

Electron Cooling by Vibrational Excitation of Carbon Dioxide

L Campbell¹, L R Hargreaves¹, P A Thorn¹, M J Brunger¹ and T N Rescigno²

¹School of Chemistry, Physics and Earth Sciences, Flinders University,
GPO Box 2100, Adelaide SA 5001, Australia.

²LBNL, Computing Sciences, Berkeley, California 94720, USA.

Abstract

We report new results for electron energy transfer rates [1] and electron cooling rates due to vibrational excitation of carbon dioxide (CO₂). The present model calculations are topical because at altitudes less than about 200 km, the neutral atmospheres of Mars and Venus are dominated by CO₂. Thus we expect electron cooling by CO₂ to be a significant energy transfer mechanism in their respective atmospheres, with vibrational excitation being the dominant process [2].

Our method was described earlier in Campbell *et al.* [1], although in this application our integral cross section (ICS) database has been meticulously constructed from the best available experimental and theoretical [e.g. 3-5] ICS for vibrational excitation in CO₂. Electron energy transfer rates from the present work are compared against those from the only previous corresponding study [2], with these results being presented at the meeting.

References

- [1] L Campbell, M J Brunger, D C Cartwright and P J O Teubner 2004, *Planetary and Space Science* **52**, 815-822.
- [2] M A Morrison and A E Greene 1978, *J Geophys Res* **83**, 1172-1174.
- [3] M J Brunger, S J Buckman and M T Elford 2003, "Photon and Electron Interactions with Atoms, Molecules and Ions", (Springer-Verlag, Berlin), 118-201.
- [4] M Allan 2001, *Phys Rev Lett* **87**, 033201.
- [5] C W McCurdy, W A Isaacs, H-D Meyer and T N Rescigno 2003, *Phys Rev A* **67**, 042708 (and references therein).